

A Property Equivalent to n -Permutability for Infinite Groups

Alireza Abdollahi* and Aliakbar Mohammadi Hassanabadi†

metadata, citation and similar papers at core.ac.uk

and

Bijan Taeri‡

Department of Mathematics, University of Technology of Isfahan, Isfahan, Iran

Communicated by Gernot Stroth

Received September 7, 1998

Let n be an integer greater than 1. A group G is said to be n -permutable whenever for every n -tuple (x_1, \dots, x_n) of elements of G there exists a non-identity permutation σ of $\{1, \dots, n\}$ such that $x_1 \cdots x_n = x_{\sigma(1)} \cdots x_{\sigma(n)}$. In this paper we prove that an infinite group G is n -permutable if and only if for every n infinite subsets X_1, \dots, X_n of G there exists a non-identity permutation σ on $\{1, \dots, n\}$ such that $X_1 \cdots X_n \cap X_{\sigma(1)} \cdots X_{\sigma(n)} \neq \emptyset$. © 1999 Academic Press

1. INTRODUCTION

Permutable groups have been studied by various people (for example, see [1–3, 5, 6]). Let n be an integer greater than 1. Recall that a group G is called n -permutable whenever for every n -tuple (x_1, \dots, x_n) of elements of G there exists a non-identity permutation σ of $\{1, \dots, n\}$ such that $x_1 \cdots x_n = x_{\sigma(1)} \cdots x_{\sigma(n)}$. Also a group is said to be permutable if it is n -permutable for some integer $n > 1$. The main result for groups in this class was obtained by Curzio *et al.* in [3], where it was shown that such

* E-mail: abdollahi@math.ui.ac.ir.

† E-mail: aamohaha@math.ui.ac.ir.

‡ E-mail: b.taeri@cc.iut.ac.ir.

groups are finite-by-abelian-by-finite. Let $n > 1$ and m be positive integers. Let S_n denote the group of all permutations on the set $\{1, \dots, n\}$. A natural extension of permutable groups, namely (m, n) -permutable groups, groups in which $X_1 \cdots X_n \subseteq \bigcup_{\sigma \in S_n \setminus 1} X_{\sigma(1)} \cdots X_{\sigma(n)}$ for all subsets X_i of G where $|X_i| = m$ for all $i = 1, \dots, n$, was introduced by Mohammadi Hassanabadi and Rhemtulla in [9]. It was proved there that such a group either is n -permutable or is finite of order bounded by a function of m and n . In [8] Mohammadi Hassanabadi investigated another extension of (m, n) -permutable groups as follows. For positive integers $n > 1$ and m a group G is called restricted (m, n) -permutable if $X_1 \cdots X_n \cap \bigcup_{\sigma \in S_n \setminus 1} X_{\sigma(1)} \cdots X_{\sigma(n)} \neq \emptyset$ for all subsets X_i of G where $|X_i| = m$ for all $i = 1, \dots, n$. It was proved there that such a group is finite-by-abelian-by-finite. In [4] Longobardi *et al.* called a group G a P_n^* -group (n an integer greater than 1) if for every sequence X_1, \dots, X_n of infinite subsets of G there exist x_i in X_i such that $x_1 \cdots x_n = x_{\sigma(1)} \cdots x_{\sigma(n)}$ for some non-trivial permutation σ in S_n . They proved that every infinite P_n^* -group is an n -permutable group. Here we deal with another extension of infinite restricted (m, n) -permutable and P_n^* -groups.

Let n be an integer greater than 1. We call a group G a restricted (∞, n) -permutable group if $X_1 \cdots X_n \cap \bigcup_{\sigma \in S_n \setminus 1} X_{\sigma(1)} \cdots X_{\sigma(n)} \neq \emptyset$ for all infinite subsets X_1, \dots, X_n of G .

Our main result is the following, which sharpens and generalizes that of [8] and also generalizes the result of [4] concerning P_n^* .

THEOREM. *Every infinite restricted (∞, n) -permutable group is n -permutable.*

2. PROOFS

To prove the theorem, we need the following results.

LEMMA 2.1. *Let G be an infinite residually finite group which is a restricted (∞, n) -permutable group. Then G is an n -permutable group.*

Proof. Let x_1, \dots, x_n be arbitrary elements of G and

$$S = \{x_1 \cdots x_n (x_{\sigma(1)} \cdots x_{\sigma(n)})^{-1} \mid \sigma \in S_n \setminus 1\}.$$

Suppose, for a contradiction, that $1 \notin S$. Since G is residually finite and S is finite, there exists a normal subgroup N of finite index in G such that $S \cap N = \emptyset$. Now considering infinite subsets Nx_1, \dots, Nx_n , there exists $\sigma \in S_n \setminus 1$ such that $Nx_1 \cdots Nx_n \cap Nx_{\sigma(1)} \cdots Nx_{\sigma(n)} \neq \emptyset$ and so $x_1 \cdots x_n (x_{\sigma(1)} \cdots x_{\sigma(n)})^{-1} \in N$, which is a contradiction. ■

LEMMA 2.2. *Let $G = Dr_{i \in I} G_i$ be an infinite direct product of non-abelian subgroups. Then G is not a restricted (∞, n) -permutable group for all integers $n > 1$.*

Proof. Suppose, for a contradiction, that G is a restricted (∞, n) -permutable group for some integer $n > 1$. We show that G is an n -permutable group, which contradicts Corollary 2.9 of [1]. Let $x_1, \dots, x_n \in G$ and put $S = \{x_1 \cdots x_n (x_{\sigma(1)} \cdots x_{\sigma(n)})^{-1} \mid \sigma \in S_n \setminus 1\}$. Let k be any integer greater than $|S|$. Since G is an infinite direct product of normal subgroups, there exist k infinite normal subgroups N_1, \dots, N_k of G such that $N_i \cap N_j = 1$ for all distinct $i, j \in \{1, \dots, k\}$. Let $l \in \{1, \dots, k\}$ and consider infinite subsets $N_l x_1, \dots, N_l x_n$. By the hypothesis, there exists $\sigma_l \in S_n \setminus 1$ such that $x_1 \cdots x_n (x_{\sigma_l(1)} \cdots x_{\sigma_l(n)})^{-1} \in N_l$. Therefore there exist two distinct $i, j \in \{1, \dots, k\}$ and an element $s \in S$ such that $s \in N_i \cap N_j = 1$ and so G is an n -permutable group. ■

We denote by A^{-1} the set $\{a^{-1} \mid a \in A\}$ for any non-empty subset A of a group. Let a and g be arbitrary elements of a group G . We define $S(a, g) := \{x \in G \mid a^x = g\}$ which is either an empty set or a right coset of the centralizer of a in G .

A key result required in the proof of the theorem is the following:

LEMMA 2.3. *Let G be an infinite restricted (∞, n) -permutable group. Then the FC-centre of G is non-trivial.*

Proof. Suppose, for a contradiction, that the FC-centre of G is trivial. We construct n infinite subsets X_1, \dots, X_n of G such that

$$X_1 \cdots X_n \cap X_{\sigma(1)} \cdots X_{\sigma(n)} = \emptyset$$

for all non-identity permutations σ in S_n . For this, for each $m \in \mathbb{N}$ we construct n subsets $X_{i,m} = \{a_{i,1}, \dots, a_{i,m}\}$ of G ($i = 1, \dots, n$), such that

$$X_{1,m} \cdots X_{n,m} \cap X_{\sigma(1),m} \cdots X_{\sigma(n),m} = \emptyset \quad (\#)$$

for all non-identity permutations σ in S_n . We argue by induction on m . Let $m = 1$. By Lemma 2.1 in [3], G is not an n -permutable group and so there exist $a_{1,1}, \dots, a_{n,1} \in G$ such that $a_{1,1} \cdots a_{n,1} \neq a_{\sigma(1),1} \cdots a_{\sigma(n),1}$ for all $\sigma \in S_n \setminus 1$. Now suppose that we have already defined subsets $X_{i,m} = \{a_{i,1}, \dots, a_{i,m}\}$ of G ($i = 1, \dots, n$) satisfying $(\#)$ for all $\sigma \in S_n \setminus 1$.

Suppose that we have already defined $a_{i,m+1}$ and so $X_{i,m+1}$ for $i = 1, \dots, r$ such that for all $\sigma \in S_n \setminus 1$

$$X_{1,m+1} X_{2,m+1} \cdots X_{r,m+1} X_{r+1,m} \cdots X_{n,m} \cap X_{\sigma(1),j_1} \cdots X_{\sigma(n),j_n} = \emptyset$$

where $j_t = m + 1$ whenever $\sigma(t) \in \{1, \dots, r\}$ and otherwise $j_t = m$. Let T_{r+1} be the union of all the following sets where σ varies over $S_n \setminus 1$,

$$X_{\sigma(i-1), s_{i-1}}^{-1} \cdots X_{\sigma(1), s_1}^{-1} X_{1, m+1} \cdots X_{r, m+1} X_{r+1, m} \cdots X_{n, m} X_{\sigma(n), s_n}^{-1} \\ \cdots X_{\sigma(i+1), s_{i+1}}^{-1}$$

where $s_l = m + 1$ whenever $\sigma(l) \in \{1, \dots, r\}$ and otherwise $s_l = m$; also i varies over $\{1, \dots, n\}$ and if $i = 1$ or $i = n$ then we define respectively $X_{\sigma(i-1), s_{i-1}}^{-1} \cdots X_{\sigma(1), s_1}^{-1} = \{1\}$ or $X_{\sigma(n), s_n}^{-1} \cdots X_{\sigma(i+1), s_{i+1}}^{-1} = \{1\}$. Also put

$$U_{r+1} = X_{r, m+1}^{-1} \cdots X_{1, m+1}^{-1} \left(\bigcup_{1 \neq \sigma \in S_n} X_{\sigma(1), j_1} \cdots X_{\sigma(n), j_n} \right) X_{n, m}^{-1} \cdots X_{r+2, m}^{-1} \\ \cup T_{r+1}$$

where $j_l = m + 1$ whenever $\sigma(l) \in \{1, \dots, r\}$ and otherwise $j_l = m$. Now we prove that there exists an element $a_{r+1, m+1} \in G \setminus U_{r+1}$ such that if $X_{r+1, m+1} = \{a_{r+1, 1}, \dots, a_{r+1, m+1}\}$ then for all $\sigma \in S_n \setminus 1$

$$X_{1, m+1} \cdots X_{r+1, m+1} X_{r+2, m} \cdots X_{n, m} \cap X_{\sigma(1), j_1} \cdots X_{\sigma(n), j_n} = \emptyset$$

where $j_l = m + 1$ whenever $\sigma(l) \in \{1, \dots, r + 1\}$ and otherwise $j_l = m$. Suppose not. Therefore $a_{1, i_1} a_{2, i_2} \cdots a_{n, i_n} = a_{\sigma(1), j_1} a_{\sigma(2), j_2} \cdots a_{\sigma(n), j_n}$ for some $1 \leq i_1, \dots, i_{r+1} \leq m + 1$, $1 \leq i_{r+2}, \dots, i_n \leq m$, $1 \leq j_1, \dots, j_n \leq m$, and $1 \leq j_s \leq m + 1$ whenever $\sigma(s) \in \{1, \dots, r + 1\}$. Suppose that $\sigma(t) = r + 1$. If $i_{r+1} \neq m + 1$ or $j_t \neq m + 1$ then we get contradiction with the induction hypothesis or the choice of $a_{r+1, m+1}$. Therefore we must always have $i_{r+1} = j_t = m + 1$ and so

$$a_{r+2, i_{r+2}} \cdots a_{n, i_n} (a_{\sigma(t+1), j_{t+1}} \cdots a_{\sigma(n), j_n})^{-1} \\ = a_{r+1, i_{r+1}}^{-1} \left(a_{1, i_1} \cdots a_{r, i_r} \right)^{-1} a_{\sigma(1), j_1} \cdots a_{\sigma(t-1), j_{t-1}} a_{r+1, j_t}.$$

Now we define g_σ and f_σ for all $\sigma \in S_n \setminus 1$ as

$$f_\sigma = \begin{cases} a_{r+2, i_{r+2}} \cdots a_{n, i_n} (a_{\sigma(t+1), j_{t+1}} \cdots a_{\sigma(n), j_n})^{-1} & \text{if } 1 \leq t \leq n - 1 \\ a_{r+2, i_{r+2}} \cdots a_{n, i_n} & \text{if } t = n \end{cases}$$

and

$$g_\sigma = \begin{cases} (a_{1, i_1} \cdots a_{r, i_r})^{-1} a_{\sigma(1), j_1} \cdots a_{\sigma(t-1), j_{t-1}} & \text{if } 2 \leq t \leq n \\ (a_{1, i_1} \cdots a_{r, i_r})^{-1} & \text{if } t = 1 \end{cases}$$

where $t = \sigma^{-1}(r + 1)$. Hence $a_{r+1, m+1} \in S(g_\sigma, f_\sigma)$ and so

$$G = U_{r+1} \cup \left(\bigcup S(g_\sigma, f_\sigma) \right) \quad (*)$$

where σ in $(*)$ varies over the set of all non-identity permutations in S_n such that $S(g_\sigma, f_\sigma) \neq \emptyset$. Obviously the set of pairs (g_σ, f_σ) is finite. Therefore $(*)$ shows that G is a finite union of right cosets of the centralizers of g_σ 's. Thus by the famous theorem of Neumann [10] there exists g_σ in the FC-centre of G such that $S(g_\sigma, f_\sigma) \neq \emptyset$. But by the hypothesis $g_\sigma = f_\sigma = 1$. Thus there exist $(n-1)$ -tuples $(i_1, \dots, i_r, i_{r+2}, \dots, i_n)$ and $(j_1, \dots, j_{t-1}, j_{t+1}, \dots, j_n)$ where $1 \leq i_1, \dots, i_r \leq m+1$, $1 \leq i_{r+2}, \dots, i_n \leq m$, $t = \sigma^{-1}(r+1)$, and $j_l = m+1$ whenever $1 \leq \sigma(l) \leq r$ and otherwise $j_l = m$ such that

$$\begin{aligned} & a_{r+2, i_{r+2}} \cdots a_{n, i_n} \left(a_{\sigma(t+1), j_{t+1}} \cdots a_{\sigma(n), j_n} \right)^{-1} \\ &= (a_{1, i_1} \cdots a_{r, i_r})^{-1} a_{\sigma(1), j_1} \cdots a_{\sigma(t-1), j_{t-1}} = 1. \end{aligned}$$

So for any $a \in X_{r+1, m}$ we have the following, which contradicts the induction hypothesis:

$$\begin{aligned} & a_{1, i_1} \cdots a_{r, i_r} a a_{r+2, i_{r+2}} \cdots a_{n, i_n} \\ &= a_{\sigma(1), j_1} a_{\sigma(2), j_2} \cdots a_{\sigma(t-1), j_{t-1}} a a_{\sigma(t+1), j_{t+1}} \cdots a_{\sigma(n), j_n}. \end{aligned}$$

Therefore we have defined $X_{r+1, m+1}$. Thus we have inductively defined $X_{i, m} = \{a_{i, 1}, \dots, a_{i, m}\}$ for all $m \in \mathbb{N}$ such that for all $\sigma \in S_n \setminus 1$

$$X_{1, m} \cdots X_{n, m} \cap X_{\sigma(1), m} \cdots X_{\sigma(n), m} = \emptyset.$$

Now set $X_i = \bigcup_{m=1}^{\infty} X_{i, m}$ ($i = 1, \dots, n$), then X_i is infinite and

$$X_1 \cdots X_n \cap X_{\sigma(1)} \cdots X_{\sigma(n)} = \emptyset$$

for all $\sigma \in S_n \setminus 1$. Otherwise there exist n -tuples (i_1, \dots, i_n) and (j_1, \dots, j_n) on \mathbb{N} and $\pi \in S_n \setminus 1$ such that $a_{1, i_1} \cdots a_{n, i_n} = a_{\pi(1), j_1} \cdots a_{\pi(n), j_n}$. Let $s = \text{Max}\{i_1, \dots, i_n, j_1, \dots, j_n\}$. Then $X_{1, s} \cdots X_{n, s} \cap X_{\pi(1), s} \cdots X_{\pi(n), s} \neq \emptyset$, which is a contradiction with the construction of $X_{i, s}$ ($i = 1, \dots, n$). ■

By Lemma 2.3, every non-trivial restricted (∞, n) -permutable group has a non-trivial FC-element and since the class of restricted (∞, n) -permutable groups is closed under homomorphic images we have:

COROLLARY 2.4. *Every restricted (∞, n) -permutable group is FC-hypercentral.*

LEMMA 2.5. *Let G be an infinite restricted (∞, n) -permutable group. If G is finitely generated or non-periodic then G is an n -permutable group.*

Proof. Suppose that G is finitely generated. By Corollary 2.4, G is FC-hypercentral. Now by a result of McLain [7] (or see p. 133 of [11]) a finitely generated FC-hypercentral group is nilpotent-by-finite. Therefore G is a finitely generated nilpotent-by-finite group and so G is residually finite. Thus G is n -permutable by Lemma 2.1. Now assume that G is non-periodic. Then there is an element x of infinite order in G . Let x_1, \dots, x_n be arbitrary elements of G . By the previous part $\langle x, x_1, \dots, x_n \rangle$ is an n -permutable group and so G is n -permutable. ■

LEMMA 2.6. *Let G be a restricted (∞, n) -permutable group. Then G is hyperabelian-by-finite.*

Proof. We may assume that G is infinite, and it suffices to show that G contains a non-trivial normal abelian subgroup. Suppose no such normal abelian subgroup exists, and let x be a non-identity element in the FC-centre of G which exists by Lemma 2.3. Let $N_1 := \langle x \rangle^G$ be the normal closure of $\langle x \rangle$ in G , and let $C := C_G(N_1)$. Then $|G : C|$ is finite and $N_1 \cap C = Z(N_1)$ is a normal abelian subgroup of G . Hence $N_1 \cap C = 1$. Therefore N_1 is finite and, having a trivial centre, it is certainly non-abelian. Now suppose, inductively, that we have already defined normal non-abelian finite subgroups N_1, \dots, N_t of G such that N_1, \dots, N_t generate their direct product in G . Write $D := C_G(N_1 \cdots N_t)$; thus $|G : D|$ is finite. Now using Lemma 2.3 we can choose a non-trivial element y in the FC-centre of D . Then y is an element of the FC-centre of G . Let $N_{t+1} := \langle y \rangle^G$. It is easily seen that N_{t+1} is a finite non-abelian group. Moreover, $N_{t+1} \subset D$, so that N_1, \dots, N_t, N_{t+1} generate their direct product in G . Thus we have found in G an infinite direct product $N_1 \times N_2 \times \cdots \times N_t \times \cdots$ of finite non-abelian groups, which together with Lemma 2.2 gives a contradiction. ■

LEMMA 2.7. *Let G be an infinite restricted (∞, n) -permutable group which is not Černikov. Then G is an n -permutable group.*

Proof. By Lemma 2.5, we may assume that G is periodic. By Lemma 2.6, there exists a normal hyperabelian subgroup H of finite index in G . Therefore H is a periodic locally soluble group and G is locally finite. Let x_1, \dots, x_n be arbitrary elements in G and let A be the finite subgroup generated by x_1, \dots, x_n . We note that H is not a Černikov group and A can be regarded as a finite group of automorphisms of H . Now by a result of Zaicev [13], there exists an abelian subgroup B of H which is not Černikov and B is a normal subgroup of AB . Since B is periodic it is a direct product of the Sylow p -subgroups B_p of B . If infinitely many B_p are non-trivial, then since $|A|$ has only finitely many prime divisors, there exists an infinite subgroup D of B which is normal in AB such that $A \cap D = 1$. Consider the n infinite subsets Dx_1, \dots, Dx_n . By the hypothesis there exists

$\sigma \in S_n \setminus 1$ such that

$$Dx_1 \cdots Dx_n \cap Dx_{\sigma(1)} \cdots Dx_{\sigma(n)} \neq \emptyset$$

and so $x_1 \cdots x_n (x_{\sigma(1)} \cdots x_{\sigma(n)})^{-1} \in A \cap D = 1$. Therefore $x_1 \cdots x_n = x_{\sigma(1)} \cdots x_{\sigma(n)}$ as required. So assume that there exist only finitely many B_p which are non-trivial. Since B is not a Černikov group and since the product of two normal Černikov subgroups of a group is a Černikov group, then there exists a prime number p such that B_p is not Černikov. Thus by Theorem 4.3.13 of [12], $C = \{b \in B \mid b^p = 1\}$ is an infinite elementary abelian p -group. Clearly C is normal in AB . Now the infinite group AC is a residually finite-by-finite group and so AC is residually finite. Therefore by Lemma 2.1, AC is an n -permutable group and the proof is complete. ■

We need the following remark in the final step of the proof of the theorem. Here $|x|$ denotes the order of an element x of a group.

REMARK 2.8. We note that if x_1, \dots, x_n ($n > 1$) are p -elements (p a prime) of distinct orders in an abelian group then $r < |x_1 \cdots x_n| \leq t$ where $r = \text{Min}\{|x_1|, \dots, |x_n|\}$ and $t = \text{Max}\{|x_1|, \dots, |x_n|\}$.

Proof of the Theorem. Let G be an infinite restricted (∞, n) -permutable group. By Lemma 2.7, we may assume that G is a Černikov group. Thus there exists an infinite normal subgroup A of G which is a direct product of finitely many groups isomorphic to C_{p^∞} , the quasicyclic p -group, for some prime number p . Let $x_1, \dots, x_n \in G$ and let X be the finite subgroup generated by x_1, \dots, x_n (we note that G is locally finite). Let Y be the group of automorphisms of A induced by the elements of X under conjugation. Then Y is finite. Let α_0 be an integer such that $|a| \leq p^{\alpha_0}$ for any $a \in X \cap A$. By Lemma 3.5 of [4] there are infinite sequences $\alpha_0 < \alpha_1 < \dots$ of integers and a_1, a_2, \dots of elements of A such that for any i , $|a_i| = p^{\alpha_i}$, and $|[a_i, y]| > p^{\alpha_{i-1}}$, for any $y \in Y \setminus C_Y(a_i)$. Now partition the set $\{a_i \mid i \geq 1\}$ into n infinite disjoint subsets J_i , $i = 1, \dots, n$. Consider the set $J_i x_i$, $i = 1, \dots, n$, and let $\sigma \in S_n \setminus 1$ be such that

$$(a_{i_1} x_1) \cdots (a_{i_n} x_n) = (a_{j_1} x_{\sigma(1)}) \cdots (a_{j_n} x_{\sigma(n)})$$

for suitable $a_{i_1} \in J_1, \dots, a_{i_n} \in J_n$ and $a_{j_1} \in J_{\sigma(1)}, \dots, a_{j_n} \in J_{\sigma(n)}$. Therefore

$$x := (x_1 \cdots x_n)^{-1} x_{\sigma(1)} \cdots x_{\sigma(n)} = a_{i_1}^{x_1 \cdots x_n} \cdots a_{i_n}^{x_n} (a_{j_1}^{x_{\sigma(1)} \cdots x_{\sigma(n)}} \cdots a_{j_n}^{x_{\sigma(n)}})^{-1}.$$

We note that i_1, \dots, i_n are pairwise distinct as are j_1, \dots, j_n . If

$$\{i_1, \dots, i_n\} \cap \{j_1, \dots, j_n\} = \emptyset$$

then by Remark 2.8, $p^{\alpha_r} < |x|$ where $r = \text{Min}\{i_1, \dots, i_n, j_1, \dots, j_n\}$, which is a contradiction, since $\alpha_r > \alpha_0$ and $x \in X \cap A$. Thus $F = \{i_1, \dots, i_n\} \cap \{j_1, \dots, j_n\} \neq \emptyset$. Let $|F| = s$. We may assume, without loss of generality, that $i_1 = j_1, \dots, i_s = j_s$. Then we may write

$$x = [a_{i_1}, y_1]^{z_1} \cdots [a_{i_s}, y_s]^{z_s} a_{i_{s+1}}^{y_{s+1}} \cdots a_{i_n}^{y_n} (a_{j_{s+1}}^{z_{s+1}} \cdots a_{j_n}^{z_n})^{-1},$$

for some $y_1, \dots, y_n, z_1, \dots, z_n \in X$. Now suppose, for a contradiction, that $x \neq 1$. If $[a_{i_1}, y_1] = \cdots = [a_{i_s}, y_s] = 1$ then $s < n$, since $x \neq 1$. Then since $i_{s+1}, \dots, i_n, j_{s+1}, \dots, j_n$ are pairwise distinct, by Remark 2.8, $|x| > p^{\alpha_k}$ where $k = \text{Min}\{i_{s+1}, \dots, i_n, j_{s+1}, \dots, j_n\}$, which is a contradiction. Thus we may assume, without loss of generality, that $y_l \in Y \setminus C_Y(a_{i_l})$, for $l = 1, \dots, s$ and $i_1 < \cdots < i_s$. Now we claim that the elements

$$[a_{i_1}, y_1], \dots, [a_{i_s}, y_s], a_{i_{s+1}}^{y_{s+1}}, \dots, a_{i_n}^{y_n}, a_{j_{s+1}}^{z_{s+1}}, \dots, a_{j_n}^{z_n}$$

have distinct orders. For, since $p^{\alpha_{i_l-1}} < |[a_{i_l}, y_l]| \leq p^{\alpha_{i_l}}$ for $l = 1, \dots, s$ and $\alpha_{i_1} < \cdots < \alpha_{i_s}$, then the elements $[a_{i_1}, y_1], \dots, [a_{i_s}, y_s]$ have distinct orders. Clearly $a_{i_{s+1}}^{y_{s+1}}, \dots, a_{i_n}^{y_n}, a_{j_{s+1}}^{z_{s+1}}, \dots, a_{j_n}^{z_n}$ have distinct orders. If there exist $l \in \{1, \dots, s\}$ and $k \in \{s+1, \dots, n\}$ such that $|[a_{i_l}, y_l]| = |a_{i_k}|$ or $|[a_{i_l}, y_l]| = |a_{j_k}|$ then since $p^{\alpha_{i_l-1}} < |[a_{i_l}, y_l]| \leq p^{\alpha_{i_l}}$, $\alpha_{i_l} = \alpha_{i_k}$ or $\alpha_{i_l} = \alpha_{j_k}$ and so $i_l = i_k$ or $i_l = j_k$, a contradiction. Now by Remark 2.8, $p^t < |x|$, where

$$p^t = \text{Min}\{|[a_{i_1}, y_1]|, \dots, |[a_{i_s}, y_s]|, |a_{i_{s+1}}^{y_{s+1}}|, \dots, |a_{i_n}^{y_n}|, |a_{j_{s+1}}^{z_{s+1}}|, \dots, |a_{j_n}^{z_n}|\}.$$

But $t > \alpha_0$ which is a contradiction. ■

REFERENCES

1. R. D. Blyth, Rewriting products of group elements, I, *J. Algebra* **116** (1988), 506–521.
2. R. D. Blyth, Rewriting products of group elements, II, *J. Algebra* **119** (1988), 246–259.
3. M. Curzio, P. Longobardi, M. Maj, and D. J. S. Robinson, A permutational property of groups, *Arch. Math.* **44** (1985), 383–389.
4. P. Longobardi, M. Maj, and A. H. Rhemtulla, Infinite groups in a given variety and Ramsey's theorem, *Comm. Algebra* **20** (1992), 127–139.
5. P. Longobardi, M. Maj, and S. E. Stonehewer, The classification of groups in which every product of four elements can be reordered, *Rend. Sem. Mat. Univ. Padova* **93** (1995), 7–26.
6. M. Maj, On the derived length of groups with some permutational property, *J. Algebra* **136** (1991), 86–91.
7. D. H. McLain, Remarks on the upper central series of a group, *Proc. Glasgow Math. Assoc.* **3** (1956), 38–44.
8. A. Mohammadi Hassanabadi, A property equivalent to permutability for groups, *Rend. Sem. Mat. Univ. Padova*, **100** (1998), 137–142.
9. A. Mohammadi Hassanabadi and A. H. Rhemtulla, Criteria for commutativity in large groups, *Canad. Math. Bull.* **41**, No. 1 (1998), 65–70.

10. B. H. Neumann, Groups covered by permutable subsets, *J. London Math. Soc.* **29** (1954), 236–248.
11. D. J. S. Robinson, “Finiteness Conditions and Generalized Soluble Groups, Part 1,” Springer-Verlag, Berlin, 1972.
12. D. J. S. Robinson, “A Course in the Theory of Groups,” 2nd ed., Springer-Verlag, Berlin, 1996.
13. D. I. Zaicev, On solvable subgroups of locally solvable groups, *Dokl. Akad. Nauk SSSR* **214** (1974), 1250–1253. [Translation in *Soviet Math. Dokl.* **15** (1974), 342–345.]